


Traumatic Bereavement in War Veterans

Anthony Papa, PhD; Yuval Neria, PhD; and Brett Litz, PhD



© 2008 Stock Marketbeat Inc. Giovanni Nguyen

More than a million men and women have been deployed to Operation Iraqi Freedom (OIF) and Operation Enduring Freedom (OEF) since the beginning of these conflicts. Recent studies show that loss of fellow service members to death or debilitating injuries is a major hazard. Hoge and colleagues reported that 68% of service members saw dead or seriously injured Americans, and 86% knew someone who was seriously injured or killed.¹ The majority of these service members will not develop lasting mental health problems as a result of the dual burden of loss and exposure to potentially traumatic events, but a salient minority will develop persistent problems.

Anthony Papa, PhD, is with the University of Nevada, Reno. Yuval Neria, PhD, is with the N.Y. State Psychiatric Institute, Columbia University College of Physicians and Surgeons, New York, New York. Brett Litz, PhD, is with the National Center for Post-Traumatic Stress Disorder, Boston University School of Medicine, Boston, Massachusetts.

Address correspondence to: Anthony Papa, PhD, University of Nevada, Reno, Department of Psychology/298, Reno, NV 89557; or e-mail apapa@unr.edu.

Dr. Papa, Dr. Neria, and Dr. Litz have disclosed no relevant financial relationships.

To date, explanations of the lasting biological and psychological sequelae of combat have been predominated by fear-conditioning models.² Unfortunately, the unique lasting impact of combat losses are overlooked or subsumed under the posttraumatic stress disorder (PTSD) construct.^{3,4} Yet in the past 10 to 15 years, clinical scientists have determined that certain losses can result in a unique and debilitating mental health syndrome, which we will refer to as complicated grief (CG).⁵ CG, which has also been named traumatic grief⁶ or prolonged grief disorder,⁷ is associated with significant suffering, functional impairment, and morbidity.^{4,5,7,8} with indications that, once individuals develop CG, they are at substantial risk for chronic problems that are resistant to treatment.⁹ Studies have repeatedly shown that CG is associated with increased frequency of PTSD, depression, functional problems,⁴ hospitalization for mental illness,¹⁰ suicidality,¹¹ medical comorbidity, such as cancer,¹² cardiac events and high blood pressure,¹³ and mortality.¹⁴ Further, our previous work with traumatic loss due to terrorism^{3,4,15} or combat in Israel^{16,17} suggests that loss from a violent death may be associated with more severe CG. In this article, we examine evidence for the existence of CG in war veterans, explore the impact this has on the mental health and fitness of these service members, and discuss the implications for treatment.

COMPLICATED GRIEF

Grief is a universal experience that involves intense emotional response to loss of a close or intimate friend or family member that is typified by sadness, longing, and anger. Normal grief is common in the first weeks and months after loss, may be intense at times, and is often accompanied by somatic, depressive, and PTSD symptoms. Yet symptoms of normal grief typically remit well within 6 months of a loss. The majority of bereaved are able to accommodate to loss and regain function-

ing.¹⁸ However, a salient minority of individuals develop persistent CG, which is manifested by chronic mourning, yearning, rumination, loneliness, loss-related anguish, and withdrawal.^{19,20}

*Simon et al found that
32% of a midlife, civilian
clinical sample that met
criteria for a CG diagnosis from
a clinical interview did not
display a comorbid MDD or
PTSD diagnosis.*

Although not included in the *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition (DSM-IV), research is accumulating to support the construct validity for a CG. Also, there is accumulating expert consensus regarding the core features for CG,^{19,20} and it is being considered for inclusion in the upcoming DSM-V. The proposed symptoms of CG include the following: 1) constant longing, yearning or pining for the lost person, 2) intrusive thoughts about the deceased, and/or 3) intense feelings of emotional pain, sorrow, or pangs of grief related to separation distress, along with five of the following; a) avoidance of reminders of the loss, b) feeling stunned, shocked, or dazed by the loss, c) confusion about role in life or a diminished sense of self, d) trouble accepting the loss, e) difficulty trusting others since the loss, f) feelings of bitterness and anger over the loss, g) difficulty moving on (eg, making new friends, pursuing new interests, h) feeling emotionally numb since the loss, and i) feeling that life is unfulfilling, empty, or meaningless without the deceased.⁷ Also in the proposed criteria, diagnosis requires persistent severe grief symptoms, which do not remit

after the first 6 months of the loss, and which interfere with functioning.⁷ Because CG is associated with considerable morbidity (eg, depression, suicidal ideation, high blood pressure), adverse health behaviors (eg, increased smoking and alcohol consumption), and quality of life impairments, the costs of unrecognized and untreated CG are high in terms of healthcare costs, lost productivity, suicide, and quality adjusted life years.^{7,8,11,13,21}

HOW COMMON IS COMPLICATED GRIEF?

Research is relatively scant on prevalence rates of CG and varies according to type of bereavement. There is no research on the prevalence rates of CG from combat. Estimates of disruption vary from 10% following deaths from natural causes,¹¹ 20% in bereaved dementia caregivers,²² to 30% to 40% in Alzheimer's caregivers^{23,24} evidencing clinically significant levels of depression. Research on the risk factors specific to CG suggests that insults to secure attachment are central to the disorder's etiology. These risk factors include childhood abuse and serious neglect,²⁵ childhood separation anxiety,²⁶ close kinship relationship to the deceased (eg, parents are the most adversely affected adults),^{27,28} insecure attachment styles,²⁸ and marital closeness, support, and dependency.²⁸⁻³⁰ Advance preparation for the loss³¹ as well as having a good social support network³² have been associated with a lower risk for CG. Taken together, these results suggest that bereaved patients with attachment difficulties, and those who feel unprepared before the death and unsupported after it are at heightened risk of CG.

COMPLICATED GRIEF COMPARED WITH DEPRESSION AND PTSD


Research is accumulating to support the construct validity for CG and its core features.^{5,15,19} In addition to unique

risk factors described above, a number of studies have found that grief symptoms (eg, yearning, searching, non-acceptance, ruminating on memories of deceased, loneliness, lack of meaning without deceased) form separate, coherent dimensions from depression symptoms (eg, fatigue, loss of energy, self-blame, depressed mood, loss of enjoyment, worthlessness) with unique outcomes in civilians^{5,15,33-36} and in combat veterans.³⁷ Furthermore, these factors are independently predictive of differential responding to psychosocial³⁵ and psychopharmacological intervention⁸ in broad-based community samples. Similar differences in constructs have been found between anxiety symptoms (eg, worry, restlessness, anxiety, nervousness) and CG.^{5,7,33,34}

Symptoms of CG and PTSD may co-occur in the event of traumatic loss,³ but avoidance of fear-inducing stimuli associated with psychic trauma does not occur with CG following a death. Rather, there is a hyperfocus on the loss and reminders of the deceased, a desire for reconnection with the deceased, and avoidance of reminders that the person is gone rather than high arousal and negative effect. Additionally, the unique separation distress symptoms of CG, such as longing and pining for the deceased, and interpersonal attachment problems of mistrust of others and difficulty forming new relationships arising from concerns about interpersonal abandonment, are not included among criteria for PTSD.^{5,19}

Simon et al found that 32% of a midlife, civilian clinical sample that met criteria for a CG diagnosis from a clinical interview did not display a comorbid MDD or PTSD diagnosis (51% did not have comorbid PTSD).^{9,38} However, they did find that increases in CG were typically associated with increased comorbidity, but when controlling for comorbidity, CG diagnosis was associated with significant, unique functional impairment.

In a similar vein, Neria and Litz tested three hypotheses closely related to this line of research among 704 bereaved friends and family members of 9/11 victims: 1) that exposure to trauma



Perhaps the single biggest indicator of the impact of combat loss on functioning is the link between survivor guilt and poor adjustment postcombat and the associated risk for increased suicide.

will increase the risk for CG among individuals who have experienced loss; 2) that CG will be common not only among family members but also among individuals who have lost friends due to 9/11 attacks; and 3) that CG will predict functional problems above and beyond PTSD and depression.³ This study indicated that exposure to trauma increased the risk for CG twofold as compared with those who experienced loss alone; that more than one-fifth of the subjects that lost a close friend in 9/11 attacks developed CG; and that CG predicted functional problems above and beyond PTSD and depression. Supporting these results, Bonanno et al found that increases in CG symptoms predicted decrements in functioning beyond PTSD

symptoms in a midlife bereaved sample and in a bereaved sample that had lost a family member in the terrorist attacks of September 11, 2001.¹⁵

These findings have important implications for populations involved in combat. Specifically, they suggest that exposure to combat trauma may exacerbate loss-related psychopathology, CG may be highly common among service members who lost close friends, and that CG has incremental validity in predicting functioning beyond PTSD.

COMPLICATED GRIEF AND THE MILITARY

The death toll among American, Iraqi, and Afghan troops is enormous. As of September 5, 2008, there were 4,152 confirmed U.S. deaths in OIF and 582 deaths in OEF reported by the Department of Defense (<http://icasualties.org/oif/default.aspx>). In addition, the death toll among Iraqi security forces (8,593) and civilians (43,209 since 2005) is very high (<http://www.icasualties.org/oef/>; These figures are based on news reports. The actual figures are likely to be much higher.)

Perhaps the most well-established indicator of the impact of combat loss on functioning is the link between survivor guilt and poor adjustment postcombat,^{39,40} and the associated risk for increased suicide.⁴¹ This is particularly salient since the phenomenon of survivor guilt shares many features with the new diagnostic criteria or CG: yearning, trouble accepting the loss, anger or bitterness about the loss, uneasiness about moving on with life, trouble feeling connected to others, and feeling that there is no meaning future because of the loss.^{39,42,43}

As previously noted, there is a dearth of population studies that have examined the prevalence of CG in the general population, and no data are available about the prevalence and impact of CG in veterans of OIF/OEF. However, research among Vietnam veterans and Israeli war veterans provides support for

the existence of CG among war veterans from previous wars. For example, Pivar and Field³⁷ found levels of grief symptoms among Vietnam combat veterans to be higher than those found in individuals who had experienced the loss of a spouse. Additionally, these symptoms were directly related to loss of fellow service members in combat, were distinct from PTSD symptoms, and were predicted by the level of attachment to unit members, level of attachment to at least one member of the unit, and the total number of individual losses experienced by each service member.

To confirm the scope and impact of loss among veterans of combat, we examined the National Vietnam Veterans Readjustment (NVVRS) database. Participants were determined to have suffered a loss if they reported having any close personal friends who were killed in or around Vietnam. In a sample of 1,636 Vietnam veterans, 1,264 reported that they knew an American killed in Vietnam, and 860 reported they lost a close friend. Those that lost a close friend reported significantly higher symptoms, which resemble the clinical picture of CG, $t(1,616) = 4.41, P < .001$, and PTSD symptoms, $t(1,630) = 6.40, P < .001$. Additionally, the risk for CG increased as the number of close friends killed increased ($r = .23, P < .001$). Among the 860 who lost a close friend, those that reported lifetime diagnosis of depression reported higher levels of CG, $t(848) = 9.46, P < .001$, and friends killed, $t(520) = 3.37, P < .01$.

In a study of Israeli war veterans from the Yom Kippur war, Neria and colleagues described the role of traumatic loss in long term adjustment problems of Israeli war veterans and prisoners of wars,⁴⁴⁻⁴⁶ suggesting that exposure to loss is strongly related to a wide range of grief-like psychopathology (eg, depressive symptoms, anger, guilt) even 2 decades after the war.^{16,17} These studies are consistent with reports among civil-

ians suggesting that loss by malicious violence is one of the most pernicious human experiences.^{4,47,48}

Military culture and the degree of cohesion of peers within units play a crucial



Because the military setting is highly relational, losing a close friend or a leader may result in grief reactions that interfere with military duties and long-term functioning.

role in functioning and well-being during and after deployment. Studies have also repeatedly documented the buffering role of positive social support during and after exposure to combat in development of mental health problems.⁴⁹ When loss occurs, its impact produces not only continued psychiatric impairment (for some), but a downward spiral of lost social resources. Because the military setting is highly relational, losing a close friend or a leader may result in grief reactions that interfere with military duties and long term functioning.¹⁶ Indeed, research indicates that the intensity and level of interrelatedness in relationships within military units have the defining characteristics of attachment relationships.⁵⁰ Thus, one might draw two conclusions from the literature reviewed in

this paper. First, that CG a critically underrecognized and unaddressed issue in service members. Second, that service members with attachment difficulties or those who feel unprepared for the loss are at heightened risk of CG especially given the violent nature of the loss, potential disruptions of unit cohesion from the need to integrate less experienced replacement personnel, and loss of important social support resources.

IMPLICATIONS FOR TREATMENT

Clearly, the awareness of the dual burden of loss and trauma on war veterans is only now beginning to be understood. There exist an urgent need to consider the unique impact of the death of close comrades on mental health, cohesion, and social support in military personnel. Although there is broad consensus that there is a strong need for empirically supported treatments for CG,⁵¹ and significant recent efforts to meet this need in civilian populations (see Shear et al, page 662), there is perhaps an equally urgent need for secondary prevention interventions for those most at risk for developing CG in general,⁵² and in war veterans, in particular, given the numbers of veterans who have participated and continue to participate in the conflicts in Iraq and Afghanistan. Currently, there is no empirically supported treatment available for war veterans available of any kind.

In this absence, the existing literature provides some general guidelines for those professionals currently treating veterans who have experienced loss. For established cases of CG in war veterans, we recommend that providers adapt existing treatments developed for civilian population such as those of Katherine Shear and colleagues⁵³ and Paul Boelen and colleagues.⁵⁴ For those at high risk for CG but have not developed CG (ie, those showing loss-related impairment that significantly disrupts functioning), we recommend focusing efforts on well

established cognitive-behavioral strategies for mood disorders to promote stable, active routines, self-care, accommodation of loss, enhanced self-efficacy, reengagement in meaningful activities, and reattachment with significant others.

FUTURE CONSIDERATIONS

It is only in the last 10 years that the unique contribution of CG in predicting functional impairment after a loss has been recognized, and only recently have the synergistic effects of the experience of both CG and PTSD been empirically explored. As a result, the development and testing of appropriate interventions has only now begun. However, the bulk of this work has been done with civilian populations. There is an exigent need to recognize the potential dual burden of loss and trauma that a large number of service members may bear as a result of military actions in Iraq and Afghanistan, and to increase efforts in developing and validating CG treatments for war veterans in order to provide mental health providers empirical guidance to address this need.

REFERENCES

- Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med*. 2004;351(1):13-22.
- Charney DS. Psychobiological mechanisms of resilience and vulnerability: implications for successful adaptation to extreme stress. *Am J Psychiatry*. 2004;161(2):195-216.
- Neria Y, Litz B. Bereavement by traumatic means: The complex synergy of trauma and grief. *J Loss Trauma*. 2004;9:73-88.
- Neria Y, Gross R, Litz B, et al. Prevalence and psychological correlates of complicated grief among bereaved adults 2.5-3.5 years after September 11th attacks. *J Trauma Stress*. 2007;20(3):251-262.
- Prigerson HG, Bierhals AJ, Kasl SV, et al. Complicated grief as a disorder distinct from bereavement related depression and anxiety: a replication study. *Am J Psychiatry*. 1996;153(11):1484-1486.
- Melhem NM, Rosales C, Karageorge J, Reynolds CF, Frank E, Shear MK. Comorbidity of axis I disorders in patients with traumatic grief. *J Clin Psychiatry*. 2001;62(11):884-887.
- Boelen PA, Prigerson HG. The influence of symptoms of prolonged grief disorder, depression, and anxiety on quality of life among bereaved adults: a prospective study. *Eur Arch Psychiatry Clin Neurosci*. 2007;257(8):444-452.
- Ott CH. The impact of complicated grief on mental and physical health at various points in the bereavement process. *Death Stud*. 2003;27(3):249-272.
- Reynolds CF, Miller MD, Pasternak RE, et al. Treatment of bereavement-related major depressive episodes in later life: a controlled study of acute and continuation treatment with nortriptyline and interpersonal psychotherapy. *Am J Psychiatry*. 1999;156(2):202-208.
- Li J, Laursen TM, Precht DH, Olsen J, Mortensen PB. Hospitalization for mental illness among parents after the death of a child. *N Engl J Med*. 2005;352(12):1190-1196.
- Latham AE, Prigerson HG. Suicidality and bereavement: complicated grief as psychiatric disorder presenting greatest risk for suicidality. *Suicide Life Threat Behav*. 2004;34(4):350-362.
- Levav I, Kohn R, Iscovich J, Abramson JH, Tsai WY, Vigdorovich D. Cancer incidence and survival following bereavement. *Am J Public Health*. 2000;90(10):1601-1607.
- Chen JH, Bierhals AJ, Prigerson HG, Kasl SV, Mazure CM, Jacobs S. Gender differences in the effects of bereavement-related psychological distress in health outcomes. *Psychol Med*. 1999;29(2):367-380.
- Li J, Precht DH, Mortensen PB, Olsen J. Mortality in parents after death of a child in Denmark: a nationwide follow-up study. *Lancet*. 2003;361(9355):363-367.
- Bonanno GA, Neria Y, Mancini A, Coifman KG, Litz B, Insel B. Is there more to complicated grief than depression and posttraumatic stress disorder? A test of incremental validity. *J Abnorm Psychol*. 2007;116(2):342-351.
- Neria Y, Solomon Z, Ginzburg K. Posttraumatic and bereavement reactions among POWs following release from captivity: the interplay of trauma and loss. In: Malkinson R, RS, Witztum E, ed. *Traumatic and Non-traumatic Loss and Bereavement: Clinical Theory and Practice*. Madison, CT: International Universities Press; 2000:91-111.
- Neria Y. Coping with tangible and intangible traumatic losses in prisoners of war. *Isr J Psychiatry Relat Sci*. 2001;38(3-4):216-225.
- Bonanno GA. Loss, trauma, and human resilience: have we underestimated the human capacity to thrive after extremely aversive events? *Am Psychol*. 2004;59(1):20-28.
- Lichtenthal WG, Cruess DG, Prigerson HG. A case for establishing complicated grief as a distinct mental disorder in DSM-V. *Clin Psychol Rev*. 2004;24(6):637-662.
- Prigerson HG, Shear MK, Jacobs SC, et al. Consensus criteria for traumatic grief. A preliminary empirical test. *Br J Psychiatry*. 1999;174:67-73.
- Silverman GK, Jacobs SC, Kasl SV, et al. Quality of life impairments associated with diagnostic criteria for traumatic grief. *Psychol Med*. 2000;30(4):857-862.
- Schulz R, Boerner K, Shear K, Zhang S, Gitlin LN. Predictors of complicated grief among dementia caregivers: a prospective study of bereavement. *Am J Geriatr Psychiatry*. 2006;14(8):650-658.
- Robinson-Whelen S, Tada Y, MacCallum RC, McGuire L, Kiecolt-Glaser JK. Long-term caregiving: what happens when it ends? *J Abnorm Psychol*. 2001;110(4):573-584.
- Schulz R, Mendelsohn AB, Haley WE, et al. End-of-life care and the effects of bereavement on family caregivers of persons with dementia. *N Engl J Med*. 2003;349(20):1936-1942.
- Silverman GK, Johnson JG, Prigerson HG. Preliminary explorations of the effects of prior trauma and loss on risk for psychiatric disorders in recently widowed people. *Isr J Psychiatry Relat Sci*. 2001;38(3-4):202-215.
- Vanderwerker LC, Jacobs SC, Parkes CM, Prigerson HG. An exploration of associations between separation anxiety in childhood and complicated grief in later life. *J Nerv Ment Dis*. 2006;194(2):121-3.
- Cleiren M, Diekstra RF, Kerkhof AJ, van der Wal J. Mode of death and kinship in bereavement: Focusing on "who" rather than "how." *Crisis*. 1994;15(1):22-36.
- van Doorn C, Kasl SV, Beery LC, Jacobs SC, Prigerson HG. The influence of marital quality and attachment styles on traumatic grief and depressive symptoms. *J Nerv Ment Dis*. 1998;186(9):566-573.
- Johnson JG, Zhang B, Greer JA, Prigerson HG. Parental control, partner dependency, and complicated grief among widowed adults in the community. *J Nerv Ment Dis*. 2007;195(1):26-30.
- Carr D, House JS, Kessler RC, Nesse RM, Sonnega J, Wortman C. Marital quality and psychological adjustment to widowhood among older adults: A longitudinal analysis. *J Gerontol B Psychol Sci Soc Sci*. 2000;55(4):S197-207.
- Barry LC, Kasl SV, Prigerson HG. Psychiatric disorders among bereaved persons: the role of perceived circumstances of death and preparedness for death. *Am J Geriatr Psychiatry*. 2002;10(4):447-457.
- Vanderwerker LC, Prigerson HG. Social support and technological connectedness as protective factors in bereavement. *J Loss Trauma*. 2004;9(1):45-57.
- Boelen PA, van den Bout J, de Keijser J. Traumatic grief as a disorder distinct from bereavement-related depression and anxiety: A replication study with bereaved mental healthcare patients. *Am J Psychiatry*. 2003;160(7):1339-1341.
- Boelen PA, van den Bout J. Complicated grief, depression, and anxiety as distinct postloss syn-

- dromes: a confirmatory factor analysis study. *Am J Psychiatry*. 2005;162(11):2175-2177.
35. Ogrodniczuk JS, Piper WE, Joyce AS, et al. Differentiating symptoms of complicated grief and depression among psychiatric outpatients. *Can J Psychiatry*. 2003;48(2):87-93.
 36. Shear KM, Jackson CT, Essock SM, Donahue SA, Felton CJ. Screening for complicated grief among Project Liberty service recipients 18 months after September 11, 2001. *Psychiatr Serv*. 2006;57(9):1291-1297.
 37. Pivar IL, Field NP. Unresolved grief in combat veterans with PTSD. *J Anxiety Disord*. 2004;18(6):745-755.
 38. Simon NM, Shear KM, Thompson EH, et al. The prevalence and correlates of psychiatric comorbidity in individuals with complicated grief. *Compr Psychiatry*. 2007;48(5):395-399.
 39. Pearn J. Traumatic stress disorders: a classification with implications for prevention and management. *Mil Med*. 2000;165(6):434-440.
 40. Henning KR, Frueh BC. Combat guilt and its relationship to PTSD symptoms. *J Clin Psychol*. 1997;53(8):801-808.
 41. Hyer L, McCranie EW, Woods MG, Boudewyns PA. Suicidal behavior among chronic Vietnam theatre veterans with PTSD. *J Clin Psychol*. 1990;46(6):713-721.
 42. Glover H. Survival guilt and the Vietnam veteran. *J Nerv Ment Dis*. 1984;172(7):393-397.
 43. Opp RE, Samson AY. Taxonomy of guilt for combat veterans. *Prof Psych*. 1989;20(3):159-165.
 44. Neria Y, Solomon Z, Dekel R. An eighteen-year follow-up study of Israeli prisoners of war and combat veterans. *J Nerv Ment Dis*. 1998;186(3):174-182.
 45. Neria Y, Solomon Z, Ginzburg K, Dekel R, Enoch D, Ohry A. Posttraumatic residues of captivity: a follow-up of Israeli ex-prisoners of war. *J Clin Psychiatry*. 2000;61(1):39-46.
 46. Solomon Z, Neria Y, Ohry A, Waysman M, Ginzburg K. PTSD among Israeli former prisoners of war and soldiers with combat stress reaction: a longitudinal study. *Am J Psychiatry*. 1994;151(4):554-559.
 47. Pfefferbaum B, Call JA, Lensgraf SJ, et al. Traumatic grief in a convenience sample of victims seeking support services after a terrorist incident. *Ann Clin Psychiatry*. 2001;13(1):19-24.
 48. Rynearson EK, McCreery JM. Bereavement after homicide: a synergism of trauma and loss. *Am J Psychiatry*. 1993;150(2):258-261.
 49. Iversen A, Nikolaou V, Greenberg N, et al. What happens to British veterans when they leave the armed forces? *Eur J Public Health*. 2005;15(2):175-184.
 50. Davidovitz R, Mikulincer M, Shaver PR, Izsak R, Popper M. Leaders as attachment figures: leaders' attachment orientations predict leadership-related mental representations and followers' performance and mental health. *J Pers Soc Psychol*. 2007;93(4):632-650.
 51. Center for the Advancement of Health. Report on bereavement and grief research. *Death Stud*. 2004;28(6):491-575.
 52. Allumbaugh DL, Hoyt WT. Effectiveness of grief counseling: a meta-analysis. *J Couns Psychol*. 1999;46:370-380.
 53. Shear K, Frank E, Houck PR, Reynolds CF. Treatment of complicated grief: a randomized controlled trial. *JAMA*. 2005;293(21):2601-2608.
 54. Boelen PA, de Keijser J, van den Hout MA, van den Bout J. Treatment of complicated grief: a comparison between cognitive-behavioral therapy and supportive counseling. *J Consult Clin Psychol*. 2007;75(2):277-284.